



Hereditary Melanoma Panel

Clinical Features: Approximately 10% of melanoma cases are attributed to hereditary predisposition [1]. Hereditary melanoma is most frequently associated with familial atypical mole melanoma syndrome (FAMMM), which tends to occur in multiple members of the same family (also known as familial melanoma) [2, 3]. Mutations in a number of genes involved in cell proliferation and melanin biosynthesis increase the risk of melanoma development. Inheritance of these genes may manifest as multiple family members with melanoma, as multiple primary melanomas in a given individual; or as a primary melanoma with onset at an early age. In general, the overall risk of melanomas in individuals who have one or more first-degree relatives with melanoma is approximately 5–12%. Other hereditary cancer syndromes, such as hereditary breast and ovarian cancer syndrome (HBOC), Li–Fraumeni syndrome (LFS), etc., may also increase the risk of melanoma [1]. The majority of the gene mutations are transmitted in an autosomal dominant fashion. The identification of individuals at risk of developing hereditary melanoma is important in order to implement strategies for reducing the burden of early disease[3].

Our Hereditary Melanoma Panel includes mutation analysis of the 12 genes listed below.

CDKN2A	CDK4	BAP1	BRCA1	BRCA2	MC1R
TP53	WRN (REC QL2)	POLE	ACD	POT1	

Hereditary Melanoma Panel genes and associated cancers

Gene	Life time risk of melanoma	Frequency of germline mutations in melanoma	Melanoma features	Cancer Syndrome	Non melanoma tumors	References
CDKN2A	By age 80, 28% in all population, 58% in Europe, 76% in the US, and 91% in Australia	20-40%	Multiple cases of melanoma in a family, early age at diagnosis, and family members with multiple primary melanomas or pancreatic cancer	FAMMM	Pancreatic cancer, neural system tumors, nonmelanoma skin cancers, uveal melanoma, and head and neck cancers, brain tumors.	[1, 4, 5]
CDK4	74.2% by age 50	rare	Similar to those with CDKN2A mutations.	FAMMM	Squamous cell carcinoma of head and neck.	[1, 3, 6, 7]
BAP1	32% of cutaneous and uveal melanoma	84% of uveal melanoma patients with metastases	Cutaneous and ocular melanoma, uveal melanoma, nevoid melanomas	NA*	Mesothelioma, renal cancer, paragangliomas, lung adenocarcinoma, and clear cell carcinoma of the kidney.	[8-12]
BRCA1	NA	rare	NA	HBOC	Skin cancer, breast and/or ovarian cancer, etc.	[13, 14]
BRCA2	Relative risk: 2.6	4.8% of ocular melanoma	Cutaneous melanoma.	HBOC	Other skin cancer, breast and/or ovarian cancer, etc.	[1, 13, 14]
MC1R	Relative risk: 2	rare	3-4 fold more likely to have thick melanomas	NA	NA	[1, 15, 16]
TP53	NA	rare	NA	LFS	Sarcomas of bone and soft tissues, carcinomas of the breast and adrenal cortex, brain tumors, and acute leukemias, etc.	[14]
WRN (RECQL2)	NA	rare	Acral lentiginous melanomas on the palms, soles or in nail beds; mucosal melanomas in the nasal cavity or esophagus.	Werner syndrome	Thyroid neoplasms, meningioma, soft tissue sarcomas, leukemia, pre-leukemic conditions and osteosarcoma/bone neoplasms.	[1, 17]
ACD	NA	0.24% in CDKN2A negative cases	NA	Telomere biology disorders	Predisposition to hematological malignancies.	[18, 19]

POLE	NA	rare	Cutaneous melanoma	NA	Colorectal cancers and adenomas	[20]
POT1	NA	<1%	NA	Telomere biology disorders	Predisposition to hematological malignancies.	[18, 21]

*NA, Not available

Test methods:

Comprehensive sequence coverage of the coding regions and splice junctions of all genes in this panel is performed. Targets of interests are enriched and prepared for sequencing using the Agilent SureSelect system. Sequencing is performed using Illumina technology and reads are aligned to the reference sequence. Variants are identified and evaluated using a custom collection of bioinformatic tools and comprehensively interpreted by our team of directors and genetic counselors. All pathogenic and likely pathogenic variants are confirmed by Sanger sequencing. The technical sensitivity of this test is estimated to be >99% for single nucleotide changes and insertions and deletions of less than 20 bp. Our CNV detection algorithm was developed and its performance determined for the sole purpose of identifying deletions and duplications within the coding region of the gene(s) tested. Partial exonic copy number changes and rearrangements of less than 400 bp may not be detected by this methodology. Regions of high homology and repetitive regions may not be analyzed. This methodology will not detect low level mosaicism, balanced translocations, inversions, or point mutations that may be responsible for the clinical phenotype. The sensitivity of our deletion/duplication assay may be reduced when DNA extracted by an outside laboratory is provided.

Hereditary Melanoma Panel (12 genes)

Sample specifications: 3 to10 cc of blood in a purple top (EDTA) tube
 Cost: \$2000
 CPT codes: 81445
 81407
 Turn-around time: 6 weeks

Note: We cannot bill insurance for this test.

Results:

Results, along with an interpretive report, are faxed to the referring physician as soon as they are completed. All abnormal results are reported by telephone or email.

For more information about our testing options, please visit our website at dnatesting.uchicago.edu or contact us at 773-834-0555.

References:

- Bonadies, D.C. and A.E. Bale, *Hereditary melanoma*. *Curr Probl Cancer*, 2011. **35**(4): p. 162-72.
- Newton Bishop, J.A., et al., *Family studies in melanoma: identification of the atypical mole syndrome (AMS) phenotype*. *Melanoma Res*, 1994. **4**(4): p. 199-206.
- Marzuka-Alcala, A., M.J. Gabree, and H. Tsao, *Melanoma susceptibility genes and risk assessment*. *Methods Mol Biol*, 2014. **1102**: p. 381-93.
- Gabree, M. and M. Seidel, *Genetic testing by cancer site: skin*. *Cancer J*, 2012. **18**(4): p. 372-80.
- Azizi, E., et al., *Familial cutaneous malignant melanoma and tumors of the nervous system. A hereditary cancer syndrome*. *Cancer*, 1995. **76**(9): p. 1571-8.
- Puntervoll, H.E., et al., *Melanoma prone families with CDK4 germline mutation: phenotypic profile and associations with MC1R variants*. *J Med Genet*, 2013. **50**(4): p. 264-70.
- Sabir, M., et al., *Novel germline CDK4 mutations in patients with head and neck cancer*. *Hered Cancer Clin Pract*, 2012. **10**(1): p. 11.
- Njauw, C.N., et al., *Germline BAP1 inactivation is preferentially associated with metastatic ocular melanoma and cutaneous-ocular melanoma families*. *PLoS One*, 2012. **7**(4): p. e35295.
- Pilarski, R., et al., *Expanding the clinical phenotype of hereditary BAP1 cancer predisposition syndrome, reporting three new cases*. *Genes Chromosomes Cancer*, 2014. **53**(2): p. 177-82.
- Harbour, J.W., et al., *Frequent mutation of BAP1 in metastasizing uveal melanomas*. *Science*, 2010. **330**(6009): p. 1410-3.
- Testa, J.R., et al., *Germline BAP1 mutations predispose to malignant mesothelioma*. *Nat Genet*, 2011. **43**(10): p. 1022-5.
- Wiesner, T., et al., *Toward an improved definition of the tumor spectrum associated with BAP1 germline mutations*. *J Clin Oncol*, 2012. **30**(32): p. e337-40.
- Ginsburg, O.M., et al., *BRCA1 and BRCA2 families and the risk of skin cancer*. *Fam Cancer*, 2010. **9**(4): p. 489-93.
- Monnerat, C., et al., *BRCA1, BRCA2, TP53, and CDKN2A germline mutations in patients with breast cancer and cutaneous melanoma*. *Fam Cancer*, 2007. **6**(4): p. 453-61.
- Cheng, L., et al., *[Molecular diagnostics and personalized medicine in malignant melanoma]*. *Zhonghua Bing Li Xue Za Zhi*, 2014. **43**(9): p. 639-43.
- Landi, M.T., et al., *MC1R, ASIP, and DNA repair in sporadic and familial melanoma in a Mediterranean population*. *J Natl Cancer Inst*, 2005. **97**(13): p. 998-1007.
- Lauper, J.M., et al., *Spectrum and risk of neoplasia in Werner syndrome: a systematic review*. *PLoS One*, 2013. **8**(4): p. e59709.
- Potrony, M., et al., *Update in genetic susceptibility in melanoma*. *Ann Transl Med*, 2015. **3**(15): p. 210.
- Aoude, L.G., et al., *Nonsense mutations in the shelterin complex genes ACD and TERF2IP in familial melanoma*. *J Natl Cancer Inst*, 2015. **107**(2).
- Aoude, L.G., et al., *POLE mutations in families predisposed to cutaneous melanoma*. *Fam Cancer*, 2015. **14**(4): p. 621-8.
- Read, J., K.A. Wadt, and N.K. Hayward, *Melanoma genetics*. *J Med Genet*, 2016. **53**(1): p. 1-14.